

# Largactil®

chlorpromazine hydrochloride

## Consumer Medicine Information (CMI)

### What is in this leaflet

This leaflet answers some common questions about Largactil.

It does not contain all the available information.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor or pharmacist has weighed the risks of you taking Largactil against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with this medicine.**

You may need to read it again.

### What Largactil is used for

The active ingredient of Largactil is chlorpromazine hydrochloride, one of a group of medicines called phenothiazines.

Largactil is used to treat a number of conditions and may be used either for a short time or for a long time.

Largactil is used to treat various problems such as severe depression or behavioural disturbances.

Largactil can also be used to treat nausea, vomiting, severe pain and unstoppable hiccups. Your doctor may have prescribed Largactil for another reason.

**Ask your doctor if you have any questions about why Largactil has been prescribed for you.**

Largactil is available only with a doctor's prescription.

### Before you take it

#### ***When you must not take it***

**Do not take Largactil if you have an allergy or any other reaction to Largactil or any of the ingredients listed at the end of this leaflet.**

**Do not take Largactil if you have any of the following:**

- circulatory, blood or bone marrow problems
- high blood pressure
- liver problems

**Do not take Largactil after the expiry date (EXP) printed on the pack.**

If you take this medicine after the expiry date has passed, it may not work as well.

**Do not take Largactil if the packaging is torn or shows signs of tampering or if the tablet, syrup or ampoule doesn't look quite right.**

**If you are not sure whether you should start taking Largactil, contact your doctor or pharmacist.**

#### ***Before you start to take it***

**Tell your doctor or pharmacist if you have allergies to:**

- any other medicines
- any other substances, such as foods, preservatives or dyes

**Tell your doctor or pharmacist if you are pregnant or intend to become pregnant.**

Your doctor or pharmacist will discuss the possible risks and benefits of using Largactil during pregnancy.

**Tell your doctor or pharmacist if you are breast-feeding or plan to breast-feed.**

Your doctor or pharmacist will discuss the possible risks and benefits of using Largactil during breastfeeding.

**Tell your doctor or pharmacist if you have or have had any medical conditions, especially the following:**

- asthma
- kidney or liver disease
- heart disease
- blood clots
- decreased thyroid activity
- low blood calcium levels
- parkinson's disease
- prostate problems
- epilepsy - fits and seizures
- glaucoma - high pressure in the eye
- muscle weakness (myasthenia gravis)
- diabetes
- chronic respiratory disorders
- dementia

**If you have not told your doctor or pharmacist about any of the above, tell them before you start taking Largactil.**

#### ***Taking other medicines***

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from**

## **your pharmacy, supermarket or health food shop.**

Some medicines and Largactil may interfere with each other. These include:

- alcohol
- benzotropine - a drug used to treat parkinson's disease
- antacid - used to treat stomach upsets
- lithium - used to treat mood disorders
- barbiturates - a group of drugs used as sedatives
- tricyclic anti depressants
- benzodiazepines - a group of drugs used to treat anxiety and muscle spasms
- medicines used to treat a fast or irregular heart beat e.g. amiodarone, quinidine, disopyramide.
- medicines that can slow your heart beat e.g. diltiazem, verapamil.
- medicines that can reduce potassium levels in the blood e.g. diuretics, laxatives.
- other medicines that can affect your heart rate e.g. methadone, pentamidine.
- anaesthetics
- opioids - a group of pain killing drugs
- antidiabetic agents
- levodopa - a drug used to treat parkinson's disease
- valproic acid - a drug used to treat epilepsy
- amphetamines
- anticoagulants
- anticholinergics
- guanethidine
- clonidine
- thiazides - diuretics (fluid tablets)
- quinidine
- MAOI - a group of drugs used to treat depression

- prochlorperazine - a drug used to treat nausea and vomiting
- desferrioxamine - a drug used to treat iron poisoning

These medicines may be affected by Largactil, or they may affect how well it works. You may need different amounts of your medicine, or you may need to use different medicines. Your doctor or pharmacist will advise you.

Your doctor and pharmacist may have more information on medicines to be careful with or to avoid while taking Largactil.

---

## **How to take it**

---

### ***How much to take***

Your doctor or pharmacist will tell you how many tablets or how much syrup you will need to take each day. This depends on your condition and whether or not you are taking any other medicines.

**Follow all directions given to you by your doctor and pharmacist carefully.**

**If you do not understand the instructions on the pack, ask your doctor or pharmacist for help.**

### ***How to take it***

**Swallow the prescribed dose of Largactil tablets whole with a full glass of water. Do not chew the tablets.**

**Swallow the prescribed dose of Largactil syrup and follow with a glass of water.**

Largactil may be given as an injection into a muscle or as a slow injection into a vein. Largactil injections should only be given by a doctor, nurse or other trained person.

### ***When to take it***

**Take Largactil at about the same time each day.**

Taking your tablets or syrup at the same time each day will have the

best effect. It will also help you remember when to take the tablets or syrup.

### ***If you forget to take it***

**If you miss a dose, simply continue to take Largactil as advised by your doctor or pharmacist. You do not need to make up for the missed dose.**

**Do not take a double dose to make up for the dose that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you are not sure what to do, ask your doctor or pharmacist. If you have trouble remembering to take your medicine, ask your pharmacist for some hints.**

### ***How long to take it***

**Continue taking Largactil until your doctor or pharmacist tells you to stop. Do not stop using it unless your doctor or pharmacist tells you to, even if you feel better.**

### ***If you take too much (overdose)***

**Immediately telephone your doctor or pharmacist or the Poisons Information Centre (telephone Australia 13 11 26, New Zealand 0800 POISON or 0800 764766), or go to accident and emergency at your nearest hospital, if you think that you or anyone else may have taken too much Largactil. Do this even if there are no signs of discomfort or poisoning.**

You may need urgent medical attention. If you take too much Largactil, you may experience:

- drowsiness
- confusion
- restlessness
- excitement
- breathing difficulties
- fast heart rate
- muscle twitching

---

## While you are using it

---

### ***Things you must do***

**Tell any other doctors, dentists, and pharmacists who are treating you that you are taking Largactil.**

**If you plan to have surgery that needs a general anaesthetic, tell your doctor or dentist that you are taking Largactil.**

**If you become pregnant while taking Largactil, tell your doctor or pharmacist.**

**You should avoid very cold or very hot environments.**

Largactil may make it difficult for your body to regulate its temperature.

**You should avoid exposure to strong sunlight while you are taking Largactil.**

If exposure to strong sunlight is unavoidable, suitable clothing, including a hat, should be worn. A sunscreen with a high level of protection should also be used.

### ***Things you must not do***

**Do not give Largactil to anyone else, even if they have the same condition as you.**

**Do not take Largactil to treat any other complaints unless your doctor or pharmacist tells you to.**

### ***Things to be careful of***

**Be careful driving or operating machinery until you know how Largactil affects you.**

Largactil may cause drowsiness in some people. Make sure you know how you react to Largactil before you drive a car, operate machinery, or do anything else that could be dangerous if you are dizzy or light-headed. If this occurs, do not drive. If you drink alcohol, dizziness or light-headedness may be worse.

---

## Side effects

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Largactil.**

Largactil helps most people, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

If you are over 65 years of age, you may have an increased chance of getting side effects.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor or pharmacist if you notice any of the following and they worry you:**

- dizziness or light headedness
- skin problems or discolouration
- agitation
- drowsiness or tiredness
- headache
- constipation
- dry mouth
- difficulty in urinating
- blurred vision

These are the more common side effects of Largactil. Mostly these are mild and short-lived.

**If any of the following happen, stop taking Largactil and tell your doctor or pharmacist immediately or go to casualty at your nearest hospital:**

- sore throat, fever or any sign of infection
- breathing difficulty
- seizures or fits
- unwanted muscle movements of the mouth, tongue, jaw, cheeks or arms and legs.
- hardness or rigidity of the muscles.
- fast heartbeat

These are very serious side effects. You may need urgent medical attention or hospitalisation.

**Other side effects not listed above may occur in some patients. Tell your doctor or pharmacist if you notice anything that is making you feel unwell.**

**Do not be alarmed by this list of possible side effects.**

You may not experience any of them.

### ***Storage***

Largactil ampoules for injection will be stored in the pharmacy or on the ward. The injection is kept in a cool dry place, protected from light.

**Keep your tablets in the pack until it is time to take them.**

If you take the tablets out of the pack they will not keep as well.

**Keep your tablets or syrup in a cool dry place. Do not store Largactil or any other medicine in the bathroom or near a sink. Do not leave it in the car on hot days or on window sills.**

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### ***Disposal***

**If your doctor or pharmacist tells you to stop taking Largactil or the tablets or syrup has passed the expiry date, ask your pharmacist what to do with any that are left over.**

---

## Product description

---

### ***What it looks like***

Largactil tablets are white film coated tablets with embossing on one side:

10mg tablet: LG / 10

25mg tablet: LG / 25

100mg tablet: LG / 100

Tablets are available in blister packs of 100.

Largactil syrup\* is a dark orange syrupy liquid with the odour of spearmint / peppermint, and is slightly fruity. Syrup is available in an amber glass 100mL bottle with a childproof cap.

Largactil for injection is a clear liquid in a 2mL glass ampoule, in a pack of 10 ampoules.

### **Ingredients**

Largactil tablets contain:

- chlorpromazine hydrochloride 10mg, 25mg, or 100mg (active ingredient)
- lactose monohydrate
- maize starch
- colloidal silicon dioxide
- magnesium stearate
- hypromellose
- polyethylene glycol
- titanium dioxide

Largactil syrup\* contains:

- chlorpromazine hydrochloride 25mg/5mL (active ingredient)
- sucrose
- caramel
- citric acid anhydrous
- polysorbate 20
- peppermint oil
- spearmint oil
- sodium citrate
- ascorbic acid
- sodium sulphite anhydrous
- sodium metabisulphite
- sodium benzoate
- fruit cup essence
- demineralised water

Largactil ampoules contain:

- chlorpromazine hydrochloride 50mg/2mL (active ingredient)
- sodium sulfite anhydrous
- sodium metabisulphite

- sodium chloride
- sodium citrate
- water for injections

### **Sponsor**

- sanofi-aventis australia pty ltd  
12-24 Talavera Road  
Macquarie Park NSW 2113
- sanofi-aventis new zealand limited  
Level 8, James & Wells Tower  
56 Cawley Street Ellerslie  
Auckland, New Zealand

Largactil® is a registered trademark.

Australian Registration Numbers:

10mg tablet AUST R 51620

25mg tablet AUST R 51618

100mg tablet AUST R 51619

25mg/5mL syrup AUST R 118128

50mg/2mL ampoule AUST R 27511

This leaflet was prepared in September 2009

\* Largactil Syrup not available in New Zealand

#60630v4.0